

Bonnor solution in five-dimensional gravity

R. Becerril* and T. Matos

Departamento de Fisica, Centro de Investigacion y de Estudios Avanzados del Instituto Politécnico Nacional, Apartado Postal 14-740, 07000 Mexico, Distrito Federal, Mexico

(Received 16 March 1989)

From Bonnor's solution of Einstein-Maxwell theory, a new solution to five-dimensional Kaluza-Klein equations which refers to a massive source carrying a magnetic and an electric dipole is constructed.

I. INTRODUCTION

As is well known, in the 1920s, Kaluza and Klein^{1,2} proposed a unified theory of gravitation and electromagnetism by assuming that the world has four spatial dimensions, one which is compactified to a circle.

Recently many-dimensional extensions of the Kaluza-Klein theory have been given in order to achieve a geometrical unification of all the fundamental interactions.³ Also, as a matter of fact, searching for exact solutions to the five-dimensional Kaluza-Klein equations has received new impetus.

In this paper we shall use a theorem given by Matos⁴ to obtain the corresponding Bonnor dipole solution for a Kaluza-Klein field. Matos's proof is highly nontrivial; we shall prove his theorem in a very simple form in Sec. II.

II. POTENTIAL FORMALISM

We shall work with a variant of the five-dimensional theory in which it is assumed to admit one non-null Killing vector field X^μ and the five-dimensional metric $\gamma_{\mu\nu}$ reads

$$\gamma_{\mu\nu} = \begin{pmatrix} I^{-1}g_{ab} + I^2 A_a A_b & I^2 A_a \\ I^2 A_b & I^2 \end{pmatrix}, \tag{1}$$

$$\mu, \nu = 1, \dots, 5, \quad a, b = 1, \dots, 4,$$

where A_a is the electromagnetic four-potential, g_{ab} is the space-time metric, and I is the scalar field. Then vacuum field equations are characterized by

$$R_{\mu\nu}^{(5)} = 0 \tag{2}$$

where $R_{\mu\nu}^{(5)}$ is the five-dimensional Ricci tensor.⁵

Stationary fields admit a second Killing vector field Y^μ with $Y^\mu Y_\mu < 0$. In this case, one can define in a covariant manner five real potentials in the form⁶

$$\begin{aligned} \mathcal{H}^{4/3} &= I^2 = X^\mu X_\mu, \quad f = -IY^\mu Y_\mu + I^{-1}(X^\mu Y_\mu)^2, \\ \psi &= -I^{-2}X_\mu Y^\mu, \quad \chi_{,\mu} = \epsilon_{\alpha\beta\gamma\delta\mu} X^\alpha Y^\beta X^\gamma Y^\delta, \\ \epsilon_{,\mu} &= \epsilon_{\alpha\beta\gamma\delta\mu} X^\alpha Y^\beta Y^\gamma Y^\delta, \end{aligned} \tag{3}$$

where f, ψ, χ, ϵ are the gravitational, electrostatic, magnetostatic, and rotational potentials, respectively, and

$\epsilon_{\alpha\beta\gamma\delta\mu}$ is the five-dimensional Levi-Civita pseudotensor.

The five-dimensional field equations (2), in function of the potentials (3), can be derived from the Lagrangian⁶

$$\begin{aligned} \mathcal{L}^{(5)} &= \frac{\rho}{2f^2} [f_{,a} f^{,a} + (\epsilon_{,a} + \psi\chi_{,a})(\epsilon^{,a} + \psi\chi^{,a})] \\ &+ \frac{\rho}{2f} (\mathcal{H}\psi_{,a}\psi^{,a} + \mathcal{H}^{-2}\chi_{,a}\chi^{,a}) + \frac{2}{3} \frac{\rho}{\mathcal{H}^2} \mathcal{H}_{,a}\mathcal{H}^{,a}, \end{aligned} \tag{4}$$

where

$$\det \gamma = -\rho^2 = \det \begin{pmatrix} \gamma_{33} & \dots \\ \dots & \gamma_{55} \end{pmatrix}.$$

The potentials (3) are analogous to the Ernst potentials (ξ, Φ) and generate a five-dimensional Riemannian potential space (see Ref. 6).

The Ernst equations for stationary Einstein-Maxwell fields outside the sources can be deduced from a variational principle with the Lagrangian

$$\mathcal{L} = \frac{\rho}{2} F^{-2} (\xi_{,a} + 2\Phi_{,a}\bar{\Phi})(\bar{\xi}^{,a} + 2\Phi\bar{\Phi}^{,a}) + 2\rho F^{-1} \Phi_{,a}\bar{\Phi}^{,a} \tag{5}$$

(see Ref. 7). For electrostatic fields $\xi = \bar{\xi}$ and $\Phi = \bar{\Phi}$ (an overbar denotes complex conjugation) then the Lagrangian is reduced to

$$\mathcal{L} = \frac{\rho}{2} F^{-2} F_{,a} F^{,a} + 2\rho F^{-1} \Phi_{,a} \Phi^{,a}. \tag{6}$$

Now we shall prove the following.⁴

Theorem 1. For each electrostatic (magnetostatic) field of the Einstein-Maxwell theory (ξ, Φ) (with $\xi = \bar{\xi}$ and $\Phi = \bar{\Phi}$) there is one corresponding field of the five-dimensional theory by the substitution

$$\begin{aligned} \mathcal{H} = \mathcal{H}_0 = \text{const}, \quad \psi &= a\Phi, \quad \chi = b\Phi, \\ \epsilon &= -c\Phi^2, \quad f = \xi + \Phi^2, \end{aligned} \tag{7a}$$

where a, b , and c are constants on the condition that

$$ab = 2c \quad \text{and} \quad \mathcal{H}^2 a^2 + \mathcal{H}^{-2} b^2 = 4. \tag{7b}$$

Proof. It can easily be checked that the substitution (7a) reduces the Lagrangian (4) to the Lagrangian (6) if the condition (7b) holds.

In the magnetic case Φ is imaginary, but theorem 1 can

be applied in virtue of the invariance of (5) by the transformation $\Phi \rightarrow i\Phi$ (Ref. 7).

III. THE BONNOR SOLUTION

As a solution of the Einstein-Maxwell equations we take the Bonnor solution;⁸ it reads

$$dS^2 = \frac{Y^2 P^2}{\Delta^3} \left[\frac{dr^2}{Z} + d\theta^2 \right] + Z \left[\frac{Y}{P} \right]^2 \sin^2 \theta d\phi^2 - \left[\frac{P}{Y} \right]^2 dt^2, \quad (8a)$$

where

$$\begin{aligned} P &= r^2 - 2mr + e^2 \cos^2 \theta, \\ \Delta &= (r - m)^2 + \sigma^2 \cos^2 \theta, \\ Y &= r^2 + e^2 \cos^2 \theta, \quad \sigma^2 = e^2 - m^2, \\ Z &= r^2 - 2mr + e^2 \end{aligned} \quad (8b)$$

in Boyer-Lindquist coordinates ($X^1 = r, X^2 = \theta, X^3 = \phi, X^4 = t$). The Ernst and the electromagnetic potentials for this solution are

$$\xi = \frac{P^2 - (2me \cos \theta)^2}{Y^2}, \quad \Phi = i \frac{2me \cos \theta}{Y}. \quad (9)$$

Using theorem 1, we can easily find the corresponding five-dimensional potentials, one arrives at

$$\begin{aligned} f &= \frac{P^2}{Y^2}, \quad \psi = -\frac{2ame \cos \theta}{Y}, \quad \chi = -\frac{2bme \cos \theta}{Y} \\ \epsilon &= -\frac{4cm^2 e^2 \cos^2 \theta}{Y^2}, \quad I = \mathcal{H}_0^{2/3}. \end{aligned} \quad (10)$$

Elements of the matrix γ are obtained using (3) and one finds

$$\begin{aligned} g_{44} &= -\frac{P^2}{Y^2}, \quad g_{34} = 0, \quad g_{33} = \frac{Y^2 Z \sin^2 \theta}{P^2}, \\ A_3 &= \frac{2bmer \sin^2 \theta}{I^3 P}, \quad A_4 = \frac{2ame \cos \theta}{Y}. \end{aligned} \quad (11)$$

The five-dimensional metric can be written as

$$dS^{(5)2} = e^{2k} dz d\bar{z} + \frac{Y^2 Z \sin^2 \theta}{I P^2} d\phi^2 - \frac{P^2}{I Y^2} dt^2 + I^2 (A_4 dt + A_3 d\phi + dy)^2, \quad (12)$$

where $z = \rho + i\xi$, $\rho = \sqrt{r^2 - 2mr + e^2} \sin \theta$, $\xi = (r - m) \cos \theta$, and y is the fifth coordinate. In order to find the function k , one has to integrate the differential equation⁹

$$2k_{,z} = \frac{1}{(\ln \rho)_{,z}} [(\ln \rho)_{,zz} + \frac{1}{4} \text{Tr} A^2], \quad (13)$$

where the matrix $A = \gamma_{,z} \gamma^{-1}$. After the integration and the choice $a = b = \sqrt{2}$, $c = \mathcal{H}_0 = 1$, the five-dimensional metric reads

$$\begin{aligned} dS^{(5)2} &= \frac{Y^2 P^2}{\Delta^3} \left[\frac{dr^2}{Z} + d\theta^2 \right] + Z \left[\frac{Y}{P} \right]^2 \sin^2 \theta d\phi^2 - \left[\frac{P}{Y} \right]^2 dt^2 \\ &+ \left[\frac{2\sqrt{2}mer \sin^2 \theta}{P} d\phi + \frac{2\sqrt{2}me \cos \theta}{Y} dt + dy \right]^2. \end{aligned} \quad (14)$$

This solution has the appearance of the extreme Bonnor form in four-dimensional Einstein-Maxwell theory. Bonnor's solution refers to a massive source carrying a magnetic dipole or (after using a known theorem¹⁰) a mass distribution carrying an electric dipole. The five-dimensional solution (14) has both an electric and a magnetic dipole and their moments are all the same $2\sqrt{2}me$. The parameters a and b are arbitrary but are parameters of electrical charge and magnetic charge, respectively; unfortunately for the time being we do not have a clear meaning of c . The parameter m is interpreted as the mass of the source. It may be easily seen that the metric (14) is asymptotically flat and for $m=0$ and $e=0$ the metric becomes flat.

*Also at Universidad Autonoma Metropolitana, Unidad Azcapotzalco. CBI-Area de Fisica. Azcapotzalco, Av. San Pablo 180. 02200 Mexico, D.F.

¹T. Kaluza, Sitzungsber. Preuss. Akad. Wiss. Phys. Math. 966 (1921).

²O. Klein, Z. Phys. 37, 895 (1926); 46, 188 (1927).

³M. J. Duff and B. E. W. Nilsson, Phys. Rep. 130, 1 (1986).

⁴T. Matos, Ann. Phys. (Leipzig) 46, 4162 (1989).

⁵D. Kramer, in *Unified Field Theory of more than 4-dimensions Including Exact Solutions*, edited by V. de Sabbata and E.

Schmutzer (World Scientific, Singapore, 1983).

⁶G. Neugebauer, Habilitationsschrift, Jena, 1969.

⁷D. Kramer, H. Stephani, M. MacCallum, and E. Herlt, *Exact Solutions of Einstein's Field Equations* (VEB Deutscher Verlag der Wissenschaften, Berlin/Cambridge University Press, Cambridge, England, 1980).

⁸W. B. Bonnor, Z. Phys. 190, 444 (1966).

⁹X. Belinsky and R. Ruffini, Phys. Lett. 89B, 195 (1980).

¹⁰W. B. Bonnor, Proc. Phys. Soc. (London) 67, 225 (1954).